




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 31/415, 31/10, 31/18</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/16227</b> <b>(43) International Publication Date:</b> 23 April 1998 (23.04.98)
<b>(21) International Application Number:</b> PCT/US97/18670 <b>(22) International Filing Date:</b> 14 October 1997 (14.10.97) <b>(30) Priority Data:</b> 60/028,494      15 October 1996 (15.10.96)      US <b>(71) Applicant (for all designated States except US):</b> G.D. SEARLE & CO. [US/US]; Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680-5110 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> SEIBERT, Karen [US/US]; 11930 Greenwalk Drive, St. Louis, MO 63146 (US). MASFERRER, Jaime [CL/US]; 1213 Blairshire, Ballwin, MO 63011 (US). GORDON, Gary, B. [US/US]; 3282 University Avenue, Highland Park, IL 60035 (US). <b>(74) Agents:</b> BULOCK, Joseph, W. et al.; G.D. Searle & Co., Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680-5110 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> METHOD OF USING CYCLOOXYGENASE-2 INHIBITORS IN THE TREATMENT AND PREVENTION OF NEOPLASIA		
<b>(57) Abstract</b>  <p>This invention relates to the use of cyclooxygenase-2 inhibitors or derivatives thereof in preventing and treating neoplasia. In particular, the invention describes the method of preventing and treating epithelial cell neoplasia in a subject, said method comprising treating the subject with a therapeutically-effective amount of a compound of Formula (I) wherein A, R<sup>2</sup> and R<sup>3</sup> are as described in the specification.</p> <div style="text-align: right; margin-top: 20px;">  <p style="text-align: right;">(I)</p> </div>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

<b>AL</b>	Albania	<b>ES</b>	Spain	<b>LS</b>	Lesotho	<b>SI</b>	Slovenia
<b>AM</b>	Armenia	<b>FI</b>	Finland	<b>LT</b>	Lithuania	<b>SK</b>	Slovakia
<b>AT</b>	Austria	<b>FR</b>	France	<b>LU</b>	Luxembourg	<b>SN</b>	Senegal
<b>AU</b>	Australia	<b>GA</b>	Gabon	<b>LV</b>	Latvia	<b>SZ</b>	Swaziland
<b>AZ</b>	Azerbaijan	<b>GB</b>	United Kingdom	<b>MC</b>	Monaco	<b>TD</b>	Chad
<b>BA</b>	Bosnia and Herzegovina	<b>GE</b>	Georgia	<b>MD</b>	Republic of Moldova	<b>TG</b>	Togo
<b>BB</b>	Barbados	<b>GH</b>	Ghana	<b>MG</b>	Madagascar	<b>TJ</b>	Tajikistan
<b>BE</b>	Belgium	<b>GN</b>	Guinea	<b>MK</b>	The former Yugoslav	<b>TM</b>	Turkmenistan
<b>BF</b>	Burkina Faso	<b>GR</b>	Greece		Republic of Macedonia	<b>TR</b>	Turkey
<b>BG</b>	Bulgaria	<b>HU</b>	Hungary	<b>ML</b>	Mali	<b>TT</b>	Trinidad and Tobago
<b>BJ</b>	Benin	<b>IE</b>	Ireland	<b>MN</b>	Mongolia	<b>UA</b>	Ukraine
<b>BR</b>	Brazil	<b>IL</b>	Israel	<b>MR</b>	Mauritania	<b>UG</b>	Uganda
<b>BY</b>	Belarus	<b>IS</b>	Iceland	<b>MW</b>	Malawi	<b>US</b>	United States of America
<b>CA</b>	Canada	<b>IT</b>	Italy	<b>MX</b>	Mexico	<b>UZ</b>	Uzbekistan
<b>CF</b>	Central African Republic	<b>JP</b>	Japan	<b>NE</b>	Niger	<b>VN</b>	Viet Nam
<b>CG</b>	Congo	<b>KE</b>	Kenya	<b>NL</b>	Netherlands	<b>YU</b>	Yugoslavia
<b>CH</b>	Switzerland	<b>KG</b>	Kyrgyzstan	<b>NO</b>	Norway	<b>ZW</b>	Zimbabwe
<b>CI</b>	Côte d'Ivoire	<b>KP</b>	Democratic People's	<b>NZ</b>	New Zealand		
<b>CM</b>	Cameroon		Republic of Korea	<b>PL</b>	Poland		
<b>CN</b>	China	<b>KR</b>	Republic of Korea	<b>PT</b>	Portugal		
<b>CU</b>	Cuba	<b>KZ</b>	Kazakstan	<b>RO</b>	Romania		
<b>CZ</b>	Czech Republic	<b>LC</b>	Saint Lucia	<b>RU</b>	Russian Federation		
<b>DE</b>	Germany	<b>LI</b>	Liechtenstein	<b>SD</b>	Sudan		
<b>DK</b>	Denmark	<b>LK</b>	Sri Lanka	<b>SE</b>	Sweden		
<b>EE</b>	Estonia	<b>LR</b>	Liberia	<b>SG</b>	Singapore		

**METHOD OF USING CYCLOOXYGENASE-2 INHIBITORS  
IN THE TREATMENT AND PREVENTION OF NEOPLASIA**

**Field of the Invention**

5

This invention is in the field of the prevention and treatment of neoplasia. More specifically, this invention relates to the use of cyclooxygenase-2 inhibitors or derivatives thereof in preventing and treating neoplasia.

10

**Background of the Invention**

Prostaglandins play a major role in the inflammation process and the inhibition of prostaglandin production, especially production of PGG<sub>2</sub>, PGH<sub>2</sub> and PGE<sub>2</sub>, has been a common target of anti-inflammatory drug discovery. However, common non-steroidal anti-inflammatory drugs (NSAID's) that are active in reducing the prostaglandin-induced pain and swelling associated with the inflammation process are also active in affecting other prostaglandin-regulated processes not associated with the inflammation process. Thus, use of high doses of most common NSAID's can produce severe side effects, including life threatening ulcers, that limit their therapeutic potential. An alternative to NSAID's is the use of corticosteroids, which also produce adverse effects, especially when long term therapy is involved.

NSAIDs have been found to prevent the production of prostaglandins by inhibiting enzymes in the human arachidonic acid/prostaglandin pathway, including the enzyme cyclooxygenase (COX). The recent discovery of an inducible enzyme associated with inflammation (named "cyclooxygenase-2 (COX-2)" or "prostaglandin G/H synthase II") provides a viable target of inhibition which more effectively reduces inflammation and produces fewer and less

drastic side effects.

Compounds which selectively inhibit cyclooxygenase-2 have been described in U.S. patents  
5 5,380,738, 5,344,991, 5,393,790, 5,434,178,  
5,474,995, 5, 510,368 and WO documents WO96/06840,  
WO96/03388, WO96/03387, WO96/25405, WO95/15316,  
WO94/15932, WO94/27980, WO95/00501, WO94/13635,  
WO94/20480, and WO94/26731.

10

Neoplastic disease states are serious and oftentimes life-threatening conditions. These neoplastic diseases, which are characterized by rapidly-proliferating cell growth, continue to be the subject of worldwide research  
15 efforts directed toward the identification of therapeutic agents which are effective in the treatment thereof. Effective therapeutic agents prolong the survivability of the patient, inhibit the rapidly-proliferating cell growth associated with the neoplasm, or effect a regression of  
20 the neoplasm. Research in this area is primarily focused toward identifying agents which would be therapeutically effective in humans and other mammals.

Recently, the presence of COX-2 has been observed in  
25 neoplastic disease. See Masanobu Oshima et al. (Cell, 87, 803-809 (1996); and Michelle Parret et al. (International Journal of Oncology, 10, 503-507 (1997)).

[Pyrazol-1-yl]benzenesulfonamides have been described  
30 as inhibitors of cyclooxygenase-2 and have shown promise in the treatment of inflammation, arthritis, and pain, with minimal side effects in pre-clinical and clinical trials. Their use for preventing colon cancer has been described in U.S. Patent No. 5,466,823. However, their use  
35 for treating colon cancer or for treating or preventing other neoplasias has not been previously described.

The present invention is directed to the use of inhibitors of cyclooxygenase-2 for the treatment and prevention of neoplasias. Conjunctive treatment of a selective cyclooxygenase-2 inhibitor with other neoplastic agents produces a synergistic effect or alternatively reduces the toxic side effects associated with chemotherapy by reducing the concentration of the side effect-causing agent needed for therapeutic efficacy.

## **Detailed Description of the Invention**

The present invention provides a method for treating or preventing a neoplasia that produces a prostaglandin in a subject in need of such treatment or prevention, the method comprises treating the subject with a therapeutically effective amount of a cyclooxygenase-2 inhibitor or derivative thereof.

The term "treatment" includes partial or total inhibition of the neoplasia growth, spreading or metastasis, as well as partial or total destruction of the neoplasia cells.

The term "prevention" includes either preventing the onset of clinically evident neoplasia altogether or preventing the onset of a preclinically evident stage of neoplasia in individuals at risk. Also intended to be encompassed by this definition is the prevention of initiation for malignant cells or to arrest or reverse the progression of premalignant cells to malignant cells. This includes prophylactic treatment of those at risk of developing the neoplasia.

The phrase "therapeutically-effective" is intended to qualify the amount of each agent which will achieve the goal of improvement in disease severity and the frequency of incidence over treatment of each agent by itself, while

avoiding adverse side effects typically associated with alternative therapies.

The term "subject" for purposes of treatment includes  
5 any human or mammal subject who has any one of the known neoplasias, and preferably is a human subject. For methods of prevention, the subject is any human or animal subject, and preferably is a human subject who is at risk for obtaining an epithelium cell-derived neoplasia. The  
10 subject may be at risk due to exposure to carcinogenic agents, being genetically predisposed to have the neoplasia, and the like.

The term "neoplasia" includes neoplasia that produce  
15 prostaglandins or express a cyclooxygenase, including both benign and cancerous tumors, growths and polyps.

In the method above, the neoplasia that produce prostaglandins include brain cancer, bone cancer,  
20 epithelial cell-derived neoplasia (epithelial carcinoma) such as basal cell carcinoma, adenocarcinoma, gastrointestinal cancer such as lip cancer, mouth cancer, esophageal cancer, small bowel cancer and stomach cancer, colon cancer, liver cancer, bladder cancer, pancreas  
25 cancer, ovary cancer, cervical cancer, lung cancer, breast cancer and skin cancer, such as squamous cell and basal cell cancers, prostate cancer, renal cell carcinoma, and other known cancers that effect epithelial cells throughout the body. Preferably, neoplasia is selected  
30 from gastrointestinal cancer, liver cancer, bladder cancer, pancreas cancer, ovary cancer, prostate cancer, cervical cancer, lung cancer, breast cancer and skin cancer, such as squamous cell and basal cell cancers. The COX-2 inhibitors can also be used to treat the fibrosis  
35 which occurs with radiation therapy. The method can be used to treat subjects having adenomatous polyps, including those with familial adenomatous polyposis (FAP).


Additionally, the method can be used to prevent polyps from forming in patients at risk of FAP.

Inhibitors of the cyclooxygenase pathway in the metabolism of arachidonic acid used in the prevention and treatment of epithelial cell derived neoplasias may inhibit enzyme activity through a variety of mechanisms. By the way of example, the inhibitors used in the methods described herein may block the enzyme activity directly by acting as a substrate for the enzyme. The use of cyclooxygenase-2 selective inhibitors is highly advantageous in that they minimize the gastric side effects that can occur with non-selective NSAID's, especially where prolonged prophylactic treatment is expected.

The term "cyclooxygenase-2 inhibitor" denotes a compound able to inhibit cyclooxygenase-2 without significant inhibition of cyclooxygenase-1. Preferably, it includes compounds which have a cyclooxygenase-2  $IC_{50}$  of less than about 0.2  $\mu M$ , and also have a selectivity ratio of cyclooxygenase-2 inhibition over cyclooxygenase-1 inhibition of at least 50, and more preferably of at least 100. Even more preferably, the compounds have a cyclooxygenase-1  $IC_{50}$  of greater than about 1  $\mu M$ , and more preferably of greater than 10  $\mu M$ . Pyrazoles can be prepared by methods described in WO95/15316, WO95/15315 and WO96/03385. Thiophene analogs can be prepared by methods described in WO 95/00501 and WO94/15932. Oxazoles can be prepared by the methods described in PCT documents WO95/00501 and WO94/27980. Isoxazoles can be prepared by the methods described in WO96/25405. . Imidazoles can be prepared by the methods described in WO96/03388 and WO96/03387. Cyclopentene

cyclooxygenase-2 inhibitors can be prepared by the methods described in U.S. Patent No. 5,344,991 and WO 95/00501. Terphenyl compounds can be prepared by the methods described in WO96/16934. Thiazole compounds can be prepared by the methods described in WO96/03392. Pyridine compounds can be prepared by the methods described in WO96/24584 and WO96/24585.

The method provided herein relates to the use of cyclooxygenase-2 inhibitors or derivatives thereof in the prevention and treatment of derived neoplasias. In the preferred embodiments, the cyclooxygenase-2 compound is selected from the compounds of Formula I

**I**

wherein A is a substituent selected from partially unsaturated or unsaturated heterocyclyl and partially unsaturated or unsaturated carbocyclic rings;

wherein R<sup>1</sup> is at least one substituent selected from heterocyclyl, cycloalkyl, cycloalkenyl and aryl, wherein R<sup>1</sup> is optionally substituted at a substitutable position with one or more radicals selected from alkyl, haloalkyl, cyano, carboxyl, alkoxy carbonyl, hydroxyl, hydroxyalkyl, haloalkoxy, amino, alkylamino, arylamino, nitro, alkoxyalkyl, alkylsulfinyl, halo, alkoxy and alkylthio;

wherein R<sup>2</sup> is methyl or amino; and

wherein R<sup>3</sup> is a radical selected from hydrido, halo, alkyl, alkenyl, alkynyl, oxo, cyano, carboxyl, cyanoalkyl, heterocyclyloxy, alkyloxy, alkylthio, alkylcarbonyl, cycloalkyl, aryl, haloalkyl,



heterocyclyl, cycloalkenyl, aralkyl,  
heterocyclylalkyl, acyl, alkylthioalkyl,  
hydroxyalkyl, alkoxycarbonyl, arylcarbonyl,  
aralkylcarbonyl, aralkenyl, alkoxyalkyl,  
5 arylthioalkyl, aryloxyalkyl, aralkylthioalkyl,  
aralkoxyalkyl, alkoxyaralkoxyalkyl,  
alkoxycarbonylalkyl, aminocarbonyl,  
aminocarbonylalkyl, alkylaminocarbonyl, N-  
arylaminocarbonyl, N-alkyl-N-arylaminocarbonyl,  
10 alkylaminocarbonylalkyl, carboxyalkyl, alkylamino, N-  
arylamino, N-aralkylamino, N-alkyl-N-aralkylamino, N-  
alkyl-N-arylamino, aminoalkyl, alkylaminoalkyl, N-  
arylaminoalkyl, N-aralkylaminoalkyl, N-alkyl-N-  
aralkylaminoalkyl, N-alkyl-N-arylaminoalkyl, aryloxy,  
15 aralkoxy, arylthio, aralkylthio, alkylsulfinyl,  
alkylsulfonyl, aminosulfonyl, alkylaminosulfonyl, N-  
arylaminosulfonyl, arylsulfonyl, N-alkyl-N-  
arylaminosulfonyl; or a pharmaceutically-acceptable  
salt thereof.

20

A preferred class of compounds which inhibit  
cyclooxygenase-2 consists of compounds of Formula I  
wherein A is selected from 5- or 6-member partially  
unsaturated heterocyclyl, 5- or 6-member unsaturated  
25 heterocyclyl, 9- or 10-member unsaturated condensed  
heterocyclyl, lower cycloalkenyl and phenyl; wherein  
R<sup>1</sup> is selected from 5- and 6-membered heterocyclyl,  
lower cycloalkyl, lower cycloalkenyl and aryl  
selected from phenyl, biphenyl and naphthyl, wherein  
30 R<sup>1</sup> is optionally substituted at a substitutable  
position with one or more radicals selected from  
lower alkyl, lower haloalkyl, cyano, carboxyl, lower  
alkoxycarbonyl, hydroxyl, lower hydroxyalkyl, lower  
haloalkoxy, amino, lower alkylamino, phenylamino,  
35 lower alkoxyalkyl, lower alkylsulfinyl, halo, lower  
alkoxy and lower alkylthio; wherein R<sup>2</sup> is methyl or  
amino; and wherein R<sup>3</sup> is a radical selected from

hydrido, oxo, cyano, carboxyl, lower alkoxycarbonyl,  
lower carboxyalkyl, lower cyanoalkyl, halo, lower  
alkyl, lower alkyloxy, lower cycloalkyl, phenyl,  
lower haloalkyl, 5- or 6-membered heterocyclyl, lower  
5 hydroxylalkyl, lower aralkyl, acyl, phenylcarbonyl,  
lower alkoxyalkyl, 5- or 6-membered heteroaryloxy,  
aminocarbonyl, lower alkylaminocarbonyl, lower  
alkylamino, lower aminoalkyl, lower alkylaminoalkyl,  
phenyloxy, and lower aralkoxy; or a pharmaceutically-  
10 acceptable salt thereof.

A more preferred class of compounds which  
inhibit cyclooxygenase-2 consists of compounds of  
Formula I wherein A is selected from oxazolyl,  
15 isoxazolyl, furyl, thienyl, dihydrofuryl, pyrrolyl,  
pyrazolyl, thiazolyl, imidazolyl, isothiazolyl,  
benzofuryl, cyclopentenyl, cyclopentadienyl, phenyl,  
and pyridyl; wherein R<sup>1</sup> is selected from pyridyl  
optionally substituted at a substitutable position  
20 with one or more methyl radicals, and phenyl  
optionally substituted at a substitutable position  
with one or more radicals selected from methyl,  
ethyl, isopropyl, butyl, tert-butyl, isobutyl,  
pentyl, hexyl, fluoromethyl, difluoromethyl,  
25 trifluoromethyl, cyano, carboxyl, methoxycarbonyl,  
ethoxycarbonyl, hydroxyl, hydroxymethyl,  
trifluoromethoxy, amino, N-methylamino, N,N-  
dimethylamino, N-ethylamino, N,N-dipropylamino, N-  
butylamino, N-methyl-N-ethylamino, phenylamino,  
30 methoxymethyl, methylsulfinyl, fluoro, chloro, bromo,  
methoxy, ethoxy, propoxy, n-butoxy, pentoxy, and  
methylthio; wherein R<sup>2</sup> is methyl or amino; and  
wherein R<sup>3</sup> is a radical selected from hydrido, oxo,  
cyano, carboxyl, methoxycarbonyl, ethoxycarbonyl,  
35 carboxypropyl, carboxymethyl, carboxyethyl,  
cyanomethyl, fluoro, chloro, bromo, methyl, ethyl,  
isopropyl, butyl, tert-butyl, isobutyl, pentyl,

hexyl, difluoromethyl, trifluoromethyl,  
pentafluoroethyl, heptafluoropropyl, difluoroethyl,  
difluoropropyl, methoxy, ethoxy, propoxy, n-butoxy,  
pentoxy, cyclohexyl, phenyl, pyridyl, thienyl,  
5 thiazolyl, oxazolyl, furyl, pyrazinyl,  
hydroxymethyl, hydroxylpropyl, benzyl, formyl,  
phenylcarbonyl, methoxymethyl, furylmethoxy,  
aminocarbonyl, N-methylaminocarbonyl, N,N-  
dimethylaminocarbonyl, N,N-dimethylamino, N-  
10 ethylamino, N,N-dipropylamino, N-butylamino, N-  
methyl-N-ethylamino, aminomethyl, N,N-  
dimethylaminomethyl, N-methyl-N-ethylaminomethyl,  
benzyloxy, and phenyloxy; or a pharmaceutically-  
acceptable salt thereof.

15

A family of specific compounds of particular  
interest within Formula I consists of compounds and  
pharmaceutically-acceptable salts thereof as  
follows:

20 5-(4-fluorophenyl)-1-[4-(methylsulfonyl)phenyl]-3-  
(trifluoromethyl)pyrazole;  
4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]-1-  
phenyl-3-(trifluoromethyl)pyrazole;  
4-(5-(4-chlorophenyl)-3-(4-methoxyphenyl)-1H-pyrazol-1-  
25 yl)benzenesulfonamide  
4-(3,5-bis(4-methylphenyl)-1H-pyrazol-1-  
yl)benzenesulfonamide;  
4-(5-(4-chlorophenyl)-3-phenyl-1H-pyrazol-1-  
yl)benzenesulfonamide;  
30 4-(3,5-bis(4-methoxyphenyl)-1H-pyrazol-1-  
yl)benzenesulfonamide;  
4-(5-(4-chlorophenyl)-3-(4-methylphenyl)-1H-pyrazol-1-  
yl)benzenesulfonamide;  
4-(5-(4-chlorophenyl)-3-(4-nitrophenyl)-1H-pyrazol-1-  
35 yl)benzenesulfonamide;  
4-(5-(4-chlorophenyl)-3-(5-chloro-2-thienyl)-1H-  
pyrazol-1-yl)benzenesulfonamide;

- 4-(4-chloro-3,5-diphenyl-1H-pyrazol-1-yl)benzenesulfonamide  
4-[5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
5 4-[5-phenyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[5-(4-fluorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[5-(4-methoxyphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
10 4-[5-(4-chlorophenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
15 4-[4-chloro-5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[3-(difluoromethyl)-5-(4-methylphenyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[3-(difluoromethyl)-5-phenyl-1H-pyrazol-1-yl]benzenesulfonamide;  
20 4-[3-(difluoromethyl)-5-(4-methoxyphenyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[3-cyano-5-(4-fluorophenyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
25 4-[3-(difluoromethyl)-5-(3-fluoro-4-methoxyphenyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[5-(3-fluoro-4-methoxyphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[4-chloro-5-phenyl-1H-pyrazol-1-yl]benzenesulfonamide;  
30 4-[5-(4-chlorophenyl)-3-(hydroxymethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[5-(4-(N,N-dimethylamino)phenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
35 5-(4-fluorophenyl)-6-[4-(methylsulfonyl)phenyl]spiro[2.4]hept-5-ene;  
4-[6-(4-fluorophenyl)spiro[2.4]hept-5-en-5-

- yl]benzenesulfonamide;  
6-(4-fluorophenyl)-7-[4-  
(methylsulfonyl)phenyl]spiro[3.4]oct-6-ene;  
5-(3-chloro-4-methoxyphenyl)-6-[4-  
5 (methylsulfonyl)phenyl]spiro[2.4]hept-5-ene;  
4-[6-(3-chloro-4-methoxyphenyl)spiro[2.4]hept-5-en-5-  
yl]benzenesulfonamide;  
5-(3,5-dichloro-4-methoxyphenyl)-6-[4-  
(methylsulfonyl)phenyl]spiro[2.4]hept-5-ene;  
10 5-(3-chloro-4-fluorophenyl)-6-[4-  
(methylsulfonyl)phenyl]spiro[2.4]hept-5-ene;  
4-[6-(3,4-dichlorophenyl)spiro[2.4]hept-5-en-5-  
yl]benzenesulfonamide;  
2-(3-chloro-4-fluorophenyl)-4-(4-fluorophenyl)-5-(4-  
15 methylsulfonylphenyl)thiazole;  
2-(2-chlorophenyl)-4-(4-fluorophenyl)-5-(4-  
methylsulfonylphenyl)thiazole;  
5-(4-fluorophenyl)-4-(4-methylsulfonylphenyl)-2-  
methylthiazole;  
20 4-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)-2-  
trifluoromethylthiazole;  
4-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)-2-(2-  
thienyl)thiazole;  
4-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)-2-  
25 benzylaminothiazole;  
4-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)-2-(1-  
propylamino)thiazole;  
2-[(3,5-dichlorophenoxy)methyl]-4-(4-fluorophenyl)-5-  
[4-(methylsulfonyl)phenyl]thiazole;  
30 5-(4-fluorophenyl)-4-(4-methylsulfonylphenyl)-2-  
trifluoromethylthiazole;  
1-methylsulfonyl-4-[1,1-dimethyl-4-(4-  
fluorophenyl)cyclopenta-2,4-dien-3-yl]benzene;  
4-[4-(4-fluorophenyl)-1,1-dimethylcyclopenta-2,4-dien-  
35 3-yl]benzenesulfonamide;  
5-(4-fluorophenyl)-6-[4-  
(methylsulfonyl)phenyl]spiro[2.4]hepta-4,6-diene;

- 4-[6-(4-fluorophenyl)spiro[2.4]hepta-4,6-dien-5-yl]benzenesulfonamide;  
6-(4-fluorophenyl)-2-methoxy-5-[4-(methylsulfonyl)phenyl]-pyridine-3-carbonitrile;  
5 2-bromo-6-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]-pyridine-3-carbonitrile;  
6-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]-2-phenyl-pyridine-3-carbonitrile;  
4-[2-(4-methylpyridin-2-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;  
10 4-[2-(5-methylpyridin-3-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;  
4-[2-(2-methylpyridin-3-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;  
15 3-[1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazol-2-yl]pyridine;  
2-[1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazol-2-yl]pyridine;  
2-methyl-4-[1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazol-2-yl]pyridine;  
20 2-methyl-6-[1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazol-2-yl]pyridine;  
4-[2-(6-methylpyridin-3-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;  
25 2-(3,4-difluorophenyl)-1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazole;  
4-[2-(4-methylphenyl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;  
2-(4-chlorophenyl)-1-[4-(methylsulfonyl)phenyl]-4-methyl-1H-imidazole;  
30 2-(4-chlorophenyl)-1-[4-(methylsulfonyl)phenyl]-4-phenyl-1H-imidazole;  
2-(4-chlorophenyl)-4-(4-fluorophenyl)-1-[4-(methylsulfonyl)phenyl]-1H-imidazole;  
35 2-(3-fluoro-4-methoxyphenyl)-1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazole;

- 1-[4-(methylsulfonyl)phenyl]-2-phenyl-4-trifluoromethyl-1H-imidazole;  
2-(4-methylphenyl)-1-[4-(methylsulfonyl)phenyl]-4-trifluoromethyl-1H-imidazole;  
5 4-[2-(3-chloro-4-methylphenyl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;  
2-(3-fluoro-5-methylphenyl)-1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazole;  
10 4-[2-(3-fluoro-5-methylphenyl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;  
2-(3-methylphenyl)-1-[4-(methylsulfonyl)phenyl]-4-trifluoromethyl-1H-imidazole;  
4-[2-(3-methylphenyl)-4-trifluoromethyl-1H-imidazol-1-yl]benzenesulfonamide;  
15 1-[4-(methylsulfonyl)phenyl]-2-(3-chlorophenyl)-4-trifluoromethyl-1H-imidazole;  
4-[2-(3-chlorophenyl)-4-trifluoromethyl-1H-imidazol-1-yl]benzenesulfonamide;  
20 4-[2-phenyl-4-trifluoromethyl-1H-imidazol-1-yl]benzenesulfonamide;  
4-[2-(4-methoxy-3-chlorophenyl)-4-trifluoromethyl-1H-imidazol-1-yl]benzenesulfonamide;  
1-allyl-4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-5-(trifluoromethyl)-1H-pyrazole;  
25 4-[1-ethyl-4-(4-fluorophenyl)-5-(trifluoromethyl)-1H-pyrazol-3-yl]benzenesulfonamide;  
N-phenyl-[4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-5-(trifluoromethyl)-1H-pyrazol-1-yl]acetamide;  
30 ethyl [4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-5-(trifluoromethyl)-1H-pyrazol-1-yl]acetate;  
4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-1-(2-phenylethyl)-1H-pyrazole;  
35 4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-1-(2-phenylethyl)-5-(trifluoromethyl)pyrazole;

- 1-ethyl-4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-5-(trifluoromethyl)-1H-pyrazole;
- 5- (4-fluorophenyl)-4-(4-methylsulfonylphenyl)-2-trifluoromethyl-1H-imidazole;
- 5 4-[4-(methylsulfonyl)phenyl]-5-(2-thiophenyl)-2-(trifluoromethyl)-1H-imidazole;
- 5-(4-fluorophenyl)-2-methoxy-4-[4-(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyridine;
- 10 2-ethoxy-5-(4-fluorophenyl)-4-[4-(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyridine;
- 5-(4-fluorophenyl)-4-[4-(methylsulfonyl)phenyl]-2-(2-propynyloxy)-6-(trifluoromethyl)pyridine;
- 2-bromo-5-(4-fluorophenyl)-4-[4-(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyridine;
- 15 4-[2-(3-chloro-4-methoxyphenyl)-4,5-difluorophenyl]benzenesulfonamide;
- 1-(4-fluorophenyl)-2-[4-(methylsulfonyl)phenyl]benzene;
- 5-difluoromethyl-4-(4-methylsulfonylphenyl)-3-phenylisoxazole;
- 20 4-[3-ethyl-5-phenylisoxazol-4-yl]benzenesulfonamide;
- 4-[5-difluoromethyl-3-phenylisoxazol-4-yl]benzenesulfonamide;
- 4-[5-hydroxymethyl-3-phenylisoxazol-4-yl]benzenesulfonamide;
- 25 4-[5-methyl-3-phenyl-isoxazol-4-yl]benzenesulfonamide;
- 1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(4-fluoro-2-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 30 1-[2-(4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 1-[2-(2,4-dichlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;
- 35 1-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;



- 1-[2-(4-methylthiophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;  
1-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;  
5 4-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;  
1-[2-(4-chlorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;  
4-[2-(4-chlorophenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;  
10 4-[2-(4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;  
4-[2-(4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;  
15 1-[2-(4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;  
1-[2-(2,3-difluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;  
4-[2-(3-fluoro-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;  
20 1-[2-(3-chloro-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;  
4-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;  
25 4-[2-(2-methylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;  
ethyl 2-[4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]oxazol-2-yl]-2-benzyl-acetate;  
2-[4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]oxazol-2-yl]acetic acid;  
30 2-(tert-butyl)-4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]oxazole;  
4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]-2-phenyloxazole;  
35 4-(4-fluorophenyl)-2-methyl-5-[4-(methylsulfonyl)phenyl]oxazole; and  
4-[5-(3-fluoro-4-methoxyphenyl)-2-trifluoromethyl-4-

oxazolyl]benzenesulfonamide.

A family of specific compounds of more particular interest within Formula I consists of compounds and pharmaceutically-acceptable salts thereof as follows:

4-[5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;

4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;

4-[5-(3-fluoro-4-methoxyphenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;

3-[1-[4-(methylsulfonyl)phenyl]-4-trifluoromethyl-1H-imidazol-2-yl]pyridine;

2-methyl-5-[1-[4-(methylsulfonyl)phenyl]-4-trifluoromethyl-1H-imidazol-2-yl]pyridine;

4-[2-(5-methylpyridin-3-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;

4-[5-methyl-3-phenylisoxazol-4-yl]benzenesulfonamide;

4-[5-hydroxymethyl-3-phenylisoxazol-4-yl]benzenesulfonamide;

[2-trifluoromethyl-5-(3,4-difluorophenyl)-4-oxazolyl]benzenesulfonamide;


4-[2-methyl-4-phenyl-5-oxazolyl]benzenesulfonamide;

and

4-[5-(3-fluoro-4-methoxyphenyl)-2-trifluoromethyl]-4-oxazolyl]benzenesulfonamide.

A subclass of cyclooxygenase-2 inhibitors is selected from compounds of WO95/15316. Preferably, the cyclooxygenase-2 inhibitor is selected from compounds of Formula II

17



wherein  $R^4$  is lower haloalkyl; wherein  $R^5$  is hydrido;  
 and wherein  $R^6$  is phenyl optionally substituted at a  
 5 substitutable position with one or more radicals selected  
 from halo, lower alkylthio, lower alkylsulfonyl, cyano,  
 nitro, lower haloalkyl, lower alkyl, hydroxyl, lower  
 alkenyl, lower hydroxyalkyl, carboxyl, lower cycloalkyl,  
 lower alkylamino, lower dialkylamino, lower  
 10 alkoxycarbonyl, aminocarbonyl, lower alkoxy, lower  
 haloalkoxy, sulfamyl, five or six membered heterocyclic  
 and amino; or a pharmaceutically-acceptable salt or  
 derivative thereof.

15 A family of specific compounds of particular interest  
 within Formula II consists of compounds, pharmaceutically-  
 acceptable salts and derivatives thereof as follows:

- 20 4-[5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;
- 4-[5-phenyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;
- 4-[5-(4-fluorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;
- 25 4-[5-(4-methoxyphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;
- 4-[5-(4-chlorophenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;
- 4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;
- 30 4-[3-(difluoromethyl)-5-(4-methylphenyl)-1H-pyrazol-1-yl]benzenesulfonamide;

4-[3-(difluoromethyl)-5-phenyl-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[3-(difluoromethyl)-5-(4-methoxyphenyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
5 4-[3-(difluoromethyl)-5-(3-fluoro-4-methoxyphenyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[5-(3-fluoro-4-methoxyphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide; and  
4-[5-(4-(N,N-dimethylamino)phenyl)-3-(trifluoromethyl)-1H-  
10 pyrazol-1-yl]benzenesulfonamide.

A family of specific compounds of more particular interest within Formula II consists of compounds and pharmaceutically-acceptable salts or derivatives thereof  
15 as follows:

4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[5-(4-chlorophenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide; and  
20 4-[5-(3-fluoro-4-methoxyphenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide.

Derivatives are intended to encompass any compounds  
25 which are structurally related to the cyclooxygenase-2 inhibitors or which possess the substantially equivalent biologic activity. By way of example, such inhibitors may include, but are not limited to, prodrugs thereof.

30 The compounds utilized in the methods of the present invention may be present in the form of free bases or pharmaceutically acceptable acid addition salts thereof. The term "pharmaceutically-acceptable salts" embraces salts commonly used to form alkali metal salts and to form  
35 addition salts of free acids or free bases. The nature of the salt is not critical, provided that it is pharmaceutically-acceptable. Suitable pharmaceutically-

acceptable acid addition salts of compounds of Formula I may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric and phosphoric acid. Appropriate organic acids may be selected from aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic and sulfonic classes of organic acids, example of which are formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, mesylic, 4-hydroxybenzoic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, benzenesulfonic, pantothenic, 2-hydroxyethanesulfonic, toluenesulfonic, sulfanilic, cyclohexylaminosulfonic, stearic, algenic,  $\beta$ -hydroxybutyric, salicylic, galactaric and galacturonic acid. Suitable pharmaceutically-acceptable base addition salts of compounds of Formula I include metallic salts made from aluminum, calcium, lithium, magnesium, potassium, sodium and zinc or organic salts made from chloroprocaine, choline, N,N'-dibenzylethylenediamine, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine. All of these salts may be prepared by conventional means from the corresponding compound of Formula I by reacting, for example, the appropriate acid or base with the compound of Formula I.

### Biological Evaluation

The efficacy of cyclooxygenase-2 inhibitors as antineoplasia agents was determined in the following models:

#### Murine Lewis lung Carcinoma Model.

35

Lewis lung carcinomas were implanted sub-cutaneously into the foot pad of male C57BL/6 mice. The mice were

subsequently treated with 4-[5-(4-chlorophenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide. The drug was supplied in the drinking water at 6 mg/kg/day. Also a non-selective COX-1/COX-2 inhibitor indomethacin was tested in this model. The drug was supplied in the drinking water at the maximum tolerated dose of 2 mg/kg/day. A total of 10 mice/compound were tested. Tumor volume was determined twice a week using a plethysmometer. The efficacy of these compounds on tumor growth was measured at day 32 after cancer cell injection, as indicated in Table 1. The % inhibition value is calculated by calculating the difference in tumor size compared with the control group.

Table 1.  
Tumor Volume (Day 32)

Treatment	% Inhibition
Vehicle/control	0.00
COX-2 inhibitor	70.86
Indomethacin	62.90

#### **Human prostate cancer cell tumors**

Two human prostate cancer cell lines (PC-3 and LNCaP) were obtained (ATCC) to determine the efficacy of cyclooxygenase-2 inhibitors to inhibit tumor growth in a therapeutic model. In addition, the LNCaP cell line secretes prostate serum antigen (PSA) when grown in nude mice.

25

PC-3

PC-3 cells ( $10^6$  cells/0.2 ml of 30% matrigel) in RPMI 1640 medium was injected on the back of nude mice. At day 28, a COX-2 inhibitor 4-[5-(4-chlorophenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide (20 mg/kg/day in water) was administered. After 45 days,  $PGE_2$

and  $\text{TXB}_2$  were measured. The COX-2 inhibitor inhibited tumor growth by 55%.  $\text{PGE}_2$  and  $\text{TXB}_2$  levels were reduced by 80-90% in the animals treated with the COX-2 inhibitor.

5 LNCaP

Similar to the results in PC-3, a COX-2 inhibitor 4-[5-(4-chlorophenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide at a dose equivalent to 6 mg/kg/day in the drinking water inhibited the growth of the tumor by 55% at day 58. PSA level was reduced to approximately 50% as judged by western blotting.

Others

Cell lines: The following cell lines can be used:  
15 classic small cell lung cancer (SCLC) cell lines NCI-H209, NCI-H345, and NCI-H510; variant SCLC cell lines NCI-N417 and NCI-H82; large cell carcinoma cell line NCI-H1155; adeno carcinoma cell line NCI-H23; and bronchioalveolar carcinoma cell line A549, breast cancer cell line MCF-7  
20 (American Type Tissue Culture Rockville MD; ATCC) and colon cancer cell lines such as NCI-H630 (ATCC), HT 29, SW948, HCA-7 and others that can be tested in vivo or in vitro. All cells can be grown in RPMI-1640, supplemented with 5% fetal bovine serum (FBS), penicillin and  
25 streptomycin (Gibco, Grand Island, NY), and be maintained in a 5%  $\text{CO}_2$  atmosphere at 37°C. All cell lines are free of mycoplasma contamination.

Growth studies: A modification (Promega CellTiter 96®, Promega Madison, WI) of the semiautomated  
30 colorimetric assay, MTT [Nakanishi, et al. *Exper. Cell Biol.*, **56**, 74-85 (1988)], which quantitates cell numbers based on reduction of a tetrazolium compound by tumor cells as determined by a spectrophotometer (540 nm) is  
35 used. All assays are performed in RPMI-1640 media supplemented with transferrin ~10 g/ml, insulin ~5 g/ml and selenium (Sigma Chemicals, St. Louis, MO). Seeding

densities are  $\sim 2 \times 10^4$  cells/well, and cells are grown for 5 days. Each experiment is reported as mean optical density corrected for background +/- standard deviation. The cyclooxygenase-2 inhibitors should be active, at a dose of 20 mg/kg, in inhibiting growth of the cancerous cell lines.

A mouse urinary bladder tumor model is performed with materials, reagents and procedures essentially as described by Grubbs et al, [*Anticancer Res.*, **13**, 33-36 (1993)]. A COX-2 inhibitor should be active at a dose of 20 mg/kg.

A rat mammary tumor model is performed with materials, reagents and procedures essentially as described by Grubbs et al., [*Anticancer Res.*, **15**, 709-16 (1995)]. A COX-2 inhibitor should be active at a dose of 20 mg/kg.

A mouse cervical and vaginal carcinogenesis model is performed with materials, reagents and procedures essentially as described by Arbeit et al., [*Proc. Acad. Sci. USA.*, **93**, 2930-35 (1996)]. A COX-2 inhibitor should be active at a dose of 20 mg/kg.

A colon adenocarcinoma cell model is performed with materials, reagents and procedures essentially as described by Shiff et al., [*J. Clin. Invest.*, **96**, 491-503 (1995)]. A COX-2 inhibitor should be active at a dose of 20 mg/kg. See also Masahiko Tsujii et al. (*Proc. Natl. Acad. Sci. USA* 94:3336-3340, 1997).

In summary, COX-2 inhibitors reduce tumor growth in several animal cancer models.



**Combination Therapy of a COX-2 inhibitor and other  
antineoplastic agents**

Lewis Lung carcinoma cells ( $2.5 \times 10^6$  cells) prepared  
5 from a brei carried in C57BL/6 mice were injected sub-  
cutaneously into the hind legs of mice. A COX-2  
inhibitor, 4-[5-(4-chlorophenyl)-3-(difluoromethyl)-1H-  
pyrazol-1-yl]benzenesulfonamide was given by gavage twice  
a week to groups of 10 mice at doses of 6 and 20 mg/kg.  
10 Cyclophosphamide (CTX) was injected to mice on days 5,7  
and 9 after the implantation of the tumor at a dose of 50  
mg/kg. Tumor volume was determined during the study.  
Animals were sacrificed at day 26 and the results of this  
experiments are summarized in Table 2. The % inhibition  
15 was calculated as above.

Table 2.

Tumor Volume (Day 22)

Treatment	% Inhibition
Vehicle	0
COX-2 inhibitor (6 mg/kg)	0
COX-2 inhibitor (20 mg/kg)	54
CTX (50 mg/kg)	57
CTX+COX-2 inhibitor (6 mg/kg)	69
CTX+COX-2 inhibitor (20 mg/kg)	77

20 The results of this experiment indicate that the  
combination of a COX-2 inhibitor and a cytotoxic agent  
produced an additive effect on their individual capacity  
to inhibit tumor growth.

25 The active compounds of the present invention may be  
administered by any suitable route known to those skilled  
in the art, preferably in the form of a pharmaceutical  
composition adapted to such a route, and in a dose

effective for the treatment intended. The active compounds and composition may, for example, be administered orally, intravascularly, intraperitoneally, intranasal, intrabronchial, subcutaneously, intra-  
5 muscularly or topically (including aerosol).

The administration of the present invention may be for either prevention or treatment purposes. The methods and compositions used herein may be used alone or in  
10 conjunction with additional therapies known to those skilled in the art in the prevention or treatment of neoplasia. Alternatively, the methods and compositions described herein may be used as conjunctive therapy. By way of example, the cyclooxygenase-2 inhibitor may be  
15 administered alone or in conjunction with other antineoplastic agents or other growth inhibiting agents or other drugs or nutrients.

There are large numbers of antineoplastic agents  
20 available in commercial use, in clinical evaluation and in pre-clinical development, which could be selected for treatment of neoplasia by combination drug chemotherapy. Such antineoplastic agents fall into several major categories, namely, antibiotic-type agents, alkylating  
25 agents, antimetabolite agents, hormonal agents, immunological agents, interferon-type agents and a category of miscellaneous agents. Alternatively, other anti-neoplastic agents, such as metallomatrix proteases (MMP), SOD mimics or  $\alpha_v\beta_3$  inhibitors may be used.

30

A first family of antineoplastic agents which may be used in combination with a selective cyclooxygenase-2 inhibitor consists of antimetabolite-type antineoplastic agents. Suitable antimetabolite antineoplastic agents may  
35 be selected from the group consisting of 5-FU-fibrinogen, acanthifolic acid, aminothiadiazone, brequinar sodium, carmofur, Ciba-Geigy CGP-30694, cyclopentyl cytosine,

cytarabine phosphate stearate, cytarabine conjugates,  
Lilly DATHF, Merrel Dow DDFC, dezaguanine,  
dideoxycytidine, dideoxyguanosine, didox, Yoshitomi DMDC,  
doxifluridine, Wellcome EHNA, Merck & Co. EX-015,  
5 fazarabine, floxuridine, fludarabine phosphate, 5-  
fluorouracil, N-(2'-furanidyl)-5-fluorouracil, Daiichi  
Seiyaku FO-152, isopropyl pyrrolizine, Lilly LY-188011,  
Lilly LY-264618, methobenzaprim, methotrexate, Wellcome  
MZPES, norspermidine, NCI NSC-127716, NCI NSC-264880, NCI  
10 NSC-39661, NCI NSC-612567, Warner-Lambert PALA,  
pentostatin, piritrexim, plicamycin, Asahi Chemical PL-AC,  
Takeda TAC-788, thioguanine, tiazofurin, Erbamont TIF,  
trimetrexate, tyrosine kinase inhibitors, tyrosine protein  
kinase inhibitors, Taiho UFT and uricytin.

15

A second family of antineoplastic agents which may be  
used in combination with a selective cyclooxygenase-2  
inhibitor consists of alkylating-type antineoplastic  
agents. Suitable alkylating-type antineoplastic agents  
20 may be selected from the group consisting of Shionogi 254-  
S, aldo-phosphamide analogues, altretamine, anaxirone,  
Boehringer Mannheim BBR-2207, bestrabucil, budotitane,  
Wakunaga CA-102, carboplatin, carmustine, Chinoin-139,  
Chinoin-153, chlorambucil, cisplatin, cyclophosphamide,  
25 American Cyanamid CL-286558, Sanofi CY-233, cyplatate,  
Degussa D-19-384, Sumimoto DACHP(Myrr)2,  
diphenylspiromustine, diplatinum cytostatic, Erba  
distamycin derivatives, Chugai DWA-2114R, ITI E09,  
elmustine, Erbamont FCE-24517, estramustine phosphate  
30 sodium, fotemustine, Unimed G-6-M, Chinoin GYKI-17230,  
hepsul-fam, ifosfamide, iproplatin, lomustine,  
mafosfamide, mitolactol, Nippon Kayaku NK-121, NCI NSC-  
264395, NCI NSC-342215, oxaliplatin, Upjohn PCNU,  
prednimustine, Proter PTT-119, ranimustine, semustine,  
35 SmithKline SK&F-101772, Yakult Honsha SN-22, spiromus-  
tine, Tanabe Seiyaku TA-077, tauromustine, temozolomide,  
teroxirone, tetraplatin and trimelamol.

A third family of antineoplastic agents which may be used in combination with a selective cyclooxygenase-2 inhibitor consists of antibiotic-type antineoplastic agents. Suitable antibiotic-type antineoplastic agents may be selected from the group consisting of Taiho 4181-A, aclarubicin, actinomycin D, actinoplanone, Erbamont ADR-456, aeroplysinin derivative, Ajinomoto AN-201-II, Ajinomoto AN-3, Nippon Soda anisomycins, anthracycline, azino-mycin-A, bisucaberin, Bristol-Myers BL-6859, Bristol-Myers BMY-25067, Bristol-Myers BMY-25551, Bristol-Myers BMY-26605, Bristol-Myers BMY-27557, Bristol-Myers BMY-28438, bleomycin sulfate, bryostatin-1, Taiho C-1027, calichemycin, chromoximycin, dactinomycin, daunorubicin, Kyowa Hakko DC-102, Kyowa Hakko DC-79, Kyowa Hakko DC-88A, Kyowa Hakko DC89-A1, Kyowa Hakko DC92-B, ditrizarubicin B, Shionogi DOB-41, doxorubicin, doxorubicin-fibrinogen, elsamicin-A, epirubicin, erbstatin, esorubicin, esperamicin-A1, esperamicin-Alb, Erbamont FCE-21954, Fujisawa FK-973, fostriecin, Fujisawa FR-900482, glidobactin, gregatin-A, grincamycin, herbimycin, idarubicin, illudins, kazusamycin, kesarirhodins, Kyowa Hakko KM-5539, Kirin Brewery KRN-8602, Kyowa Hakko KT-5432, Kyowa Hakko KT-5594, Kyowa Hakko KT-6149, American Cyanamid LL-D49194, Meiji Seika ME 2303, menogaril, mitomycin, mitoxantrone, SmithKline M-TAG, neoenactin, Nippon Kayaku NK-313, Nippon Kayaku NKT-01, SRI International NSC-357704, oxalysine, oxaunomycin, peplomycin, pilatin, pirarubicin, porothramycin, pyrindamycin A, Tobishi RA-I, rapamycin, rhizoxin, rodorubicin, sibanomicin, siwenmycin, Sumitomo SM-5887, Snow Brand SN-706, Snow Brand SN-07, sorangicin-A, sparsomycin, SS Pharmaceutical SS-21020, SS Pharmaceutical SS-7313B, SS Pharmaceutical SS-9816B, steffimycin B, Taiho 4181-2, talisomycin, Takeda TAN-868A, terpentecin, thiazine, tricrozarin A, Upjohn U-73975, Kyowa Hakko UCN-10028A, Fujisawa WF-3405, Yoshitomi Y-25024 and zorubicin.

A fourth family of antineoplastic agents which may be used in combination with the selective cyclooxygenase-2 inhibitor consists of a miscellaneous family of antineoplastic agents selected from the group consisting of alpha-carotene, alpha-difluoromethyl-arginine, acitretin, Biotec AD-5, Kyorin AHC-52, alstonine, amonafide, amphetamine, amsacrine, Angiostat, ankinomycin, anti-neoplaston A10, antineoplaston A2, antineoplaston A3, antineoplaston A5, antineoplaston AS2-1, Henkel APD, aphidicolin glycinate, asparaginase, Avarol, baccharin, batracylin, benfluron, benzotript, Ipsen-Beaufour BIM-23015, bisantrene, Bristo-Myers BMY-40481, Vestar boron-10, bromofosfamide, Wellcome BW-502, Wellcome BW-773, caracemide, carmethizole hydrochloride, Ajinomoto CDAF, chlorsulfaquinoxalone, Chemes CHX-2053, Chemex CHX-100, Warner-Lambert CI-921, Warner-Lambert CI-937, Warner-Lambert CI-941, Warner-Lambert CI-958, clanfenur, claviridenone, ICN compound 1259, ICN compound 4711, Contracan, Yakult Honsha CPT-11, crisnatol, curaderm, cytochalasin B, cytarabine, cytocytin, Merz D-609, DABIS maleate, dacarbazine, datelliptinium, didemnin-B, dihaematoporphyrin ether, dihydrolenperone, dinaline, distamycin, Toyo Pharmar DM-341, Toyo Pharmar DM-75, Daiichi Seiyaku DN-9693, elliprabin, elliptinium acetate, Tsumura EPMTc, ergotamine, etoposide, etretinate, fenretinide, Fujisawa FR-57704, gallium nitrate, genkwadaphnin, Chugai GLA-43, Glaxo GR-63178, grifolan NMF-5N, hexadecylphosphocholine, Green Cross HO-221, homoharringtonine, hydroxyurea, BTG ICRF-187, ilmofosine, isoglutamine, isotretinoin, Otsuka JI-36, Ramot K-477, Otsuka K-76COONa, Kureha Chemical K-AM, MECT Corp KI-8110, American Cyanamid L-623, leukoregulin, lonidamine, Lundbeck LU-23-112, Lilly LY-186641, NCI (US) MAP, marylzin, Merrel Dow MDL-27048, Medco MEDR-340, merbarone, merocyanine derivatives, methylanilinoacridine, Molecular

Genetics MGI-136, minactivin, mitonafide, mitoquidone, mopidamol, motretinide, Zenyaku Kogyo MST-16, N-(retinoyl)amino acids, Nisshin Flour Milling N-021, N-acylated-dehydroalanines, nafazatrom, Taisho NCU-190, 5 nocodazole derivative, Normosang, NCI NSC-145813, NCI NSC-361456, NCI NSC-604782, NCI NSC-95580, octreotide, Ono ONO-112, oquizanocine, Akzo Org-10172, pancratistatin, pazelliptine, Warner-Lambert PD-111707, Warner-Lambert PD-115934, Warner-Lambert PD-131141, Pierre Fabre PE-1001, 10 ICRT peptide D, piroxantrone, polyhaematoporphyrin, polypreic acid, Efamol porphyrin, probimane, procarbazine, proglumide, Invitron protease nexin I, Tobishi RA-700, razoxane, Sapporo Breweries RBS, restrictin-P, retelliptine, retinoic acid, Rhone-Poulenc RP-49532, 15 Rhone-Poulenc RP-56976, SmithKline SK&F-104864, Sumitomo SM-108, Kuraray SMANCS, SeaPharm SP-10094, spatol, spirocyclopropane derivatives, spirogermanium, Unimed, SS Pharmaceutical SS-554, strypoldinone, Stypoldione, Suntory SUN 0237, Suntory SUN 2071, superoxide dismutase, Toyama 20 T-506, Toyama T-680, taxol, Teijin TEI-0303, teniposide, thaliblastine, Eastman Kodak TJB-29, tocotrienol, Topostin, Teijin TT-82, Kyowa Hakko UCN-01, Kyowa Hakko UCN-1028, ukrain, Eastman Kodak USB-006, vinblastine sulfate, vincristine, vindesine, vinestramide, 25 vinorelbine, vintriptol, vinzolidine, withanolides and Yamanouchi YM-534.

Examples of radioprotective agents which may be used in the combination chemotherapy of this invention are AD- 30 5, adchnon, amifostine analogues, detox, dimesna, 1-102, MM-159, N-acylated-dehydroalanines, TGF- Genentech, tiprotimod, amifostine, WR-151327, FUT-187, ketoprofen transdermal, nabumetone, superoxide dismutase (Chiron) and superoxide dismutase Enzon.

35

Methods for preparation of the antineoplastic agents described above may be found in the literature. Methods

for preparation of doxorubicin, for example, are described in U.S. Patents No. 3,590,028 and No. 4,012,448. Methods for preparing metallomatrix protease inhibitors are described in EP 780386. Methods for preparing SOD mimics  
5 are described in EP 524,101. Methods for preparing  $\alpha,\beta$ , inhibitors are described in WO97/08174.

The phrase "conjunctive therapy" (or "combination therapy"), in defining use of a cyclooxygenase-2 inhibitor  
10 agent and another pharmaceutical agent, is intended to embrace administration of each agent in a sequential manner in a regimen that will provide beneficial effects of the drug combination, and is intended as well to embrace co-administration of these agents in a  
15 substantially simultaneous manner, such as in a single formulation having a fixed ratio of these active agents, or in multiple, separate formulations for each agent. The present invention also comprises a pharmaceutical composition for the prevention and treatment of neoplasia,  
20 comprising a therapeutically-effective amount of a compound of Formula I in association with at least one pharmaceutically-acceptable carrier, adjuvant or diluent (collectively referred to herein as "carrier" materials) and, other antineoplastic agents or other growth  
25 inhibiting agents or other drugs or nutrients.

For oral administration, the pharmaceutical composition may be in the form of, for example, a tablet, capsule, suspension or liquid. The pharmaceutical  
30 composition is preferably made in the form of a dosage unit containing a particular amount of the active ingredient. Examples of such dosage units are capsules, tablets, powders, granules or a suspension, with conventional additives such as lactose, mannitol, corn  
35 starch or potato starch; with binders such as crystalline cellulose, cellulose derivatives, acacia, corn starch or gelatins; with disintegrators such as corn starch, potato

starch or sodium carboxymethyl-cellulose; and with lubricants such as talc or magnesium stearate. The active ingredient may also be administered by injection as a composition wherein, for example, saline, dextrose or  
5 water may be used as a suitable carrier.

For intravenous, intramuscular, subcutaneous, or intraperitoneal administration, the compound may be combined with a sterile aqueous solution which is  
10 preferably isotonic with the blood of the recipient. Such formulations may be prepared by dissolving solid active ingredient in water containing physiologically compatible substances such as sodium chloride, glycine, and the like, and having a buffered pH compatible with physiological  
15 conditions to produce an aqueous solution, and rendering said solution sterile. The formulations may be present in unit or multi-dose containers such as sealed ampoules or vials.

20 If the neoplasia is localized in the G.I. tract, the compound may be formulated with acid-stable, base-labile coatings known in the art which begin to dissolve in the high pH small intestine. Formulation to enhance local pharmacologic effects and reduce systemic uptake are  
25 preferred.

Formulations suitable for parenteral administration conveniently comprise a sterile aqueous preparation of the active compound which is preferably made isotonic.  
30 Preparations for injections may also be formulated by suspending or emulsifying the compounds in non-aqueous solvent, such as vegetable oil, synthetic aliphatic acid glycerides, esters of higher aliphatic acids or propylene glycol.

35

Formulations for topical use include known gels, creams, oils, and the like. For aerosol delivery, the



compounds may be formulated with known aerosol excipients, such as saline, and administered using commercially available nebulizers. Formulation in a fatty acid source may be used to enhance biocompatibility. Aerosol delivery is the preferred method of delivery for epithelial neoplasias of the lung for prevention application.

For rectal administration, the active ingredient may be formulated into suppositories using bases which are solid at room temperature and melt or dissolve at body temperature. Commonly used bases include cocoa butter, glycerinated gelatin, hydrogenated vegetable oil, polyethylene glycols of various molecular weights, and fatty esters of polyethylene stearate.

The dosage form and amount can be readily established by reference to known neoplasia treatment or prophylactic regimens. The amount of therapeutically active compound that is administered and the dosage regimen for treating a disease condition with the compounds and/or compositions of this invention depends on a variety of factors, including the age, weight, sex and medical condition of the subject, the severity of the disease, the route and frequency of administration, and the particular compound employed, the location of the neoplasia, as well as the pharmacokinetic properties of the individual treated, and thus may vary widely. The dosage will generally be lower if the compounds are administered locally rather than systemically, and for prevention rather than for treatment. Such treatments may be administered as often as necessary and for the period of time judged necessary by the treating physician. One of skill in the art will appreciate that the dosage regime or therapeutically effective amount of the inhibitor to be administered may need to be optimized for each individual. The pharmaceutical compositions may contain active ingredient in the range of about 0.1 to 2000 mg, preferably in the


range of about 0.5 to 500 mg and most preferably between about 1 and 200 mg. A daily dose of about 0.01 to 100 mg/kg body weight, preferably between about 0.1 and about 50 mg/kg body weight, may be appropriate. The daily dose  
5 can be administered in one to four doses per day.

All patents and documents referenced herein are incorporated by reference.

10 Although this invention has been described with respect to specific embodiments, the details of these embodiments are not to be construed as limitations.

What is claimed is:

1. A method of treating a neoplasia in a subject,  
said method comprising treating the subject with a  
5 therapeutically-effective amount of a compound of Formula  
II



- 10 wherein R<sup>2</sup> is lower haloalkyl; wherein R<sup>3</sup> is hydrido;  
and wherein R<sup>4</sup> is phenyl optionally substituted at a  
substitutable position with one or more radicals selected  
from halo, lower alkylthio, lower alkylsulfonyl, cyano,  
nitro, lower haloalkyl, lower alkyl, hydroxyl, lower  
15 alkenyl, lower hydroxyalkyl, carboxyl, lower cycloalkyl,  
lower alkylamino, lower dialkylamino, lower  
alkoxycarbonyl, aminocarbonyl, lower alkoxy, lower  
haloalkoxy, sulfamyl, five or six membered heterocyclic  
and amino; or a pharmaceutically-acceptable salt or  
20 derivative thereof.

2. The method of Claim 1 wherein the compound is  
selected from compounds, and their pharmaceutically  
acceptable salts, of the group consisting of  
25 4-[5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-  
yl]benzenesulfonamide;  
4-[5-phenyl-3-(trifluoromethyl)-1H-pyrazol-1-  
yl]benzenesulfonamide;  
4-[5-(4-fluorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-  
30 yl]benzenesulfonamide;  
4-[5-(4-methoxyphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-  
yl]benzenesulfonamide;

- 4-[5-(4-chlorophenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
5 4-[3-(difluoromethyl)-5-(4-methylphenyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[3-(difluoromethyl)-5-phenyl-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[3-(difluoromethyl)-5-(4-methoxyphenyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
10 4-[3-(difluoromethyl)-5-(3-fluoro-4-methoxyphenyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[5-(3-fluoro-4-methoxyphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide; and  
15 4-[5-(4-(N,N-dimethylamino)phenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide.


3. The method of Claim 2 wherein the compound is 4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide, or a pharmaceutically-acceptable salt thereof.

4. The method of Claim 2 wherein the compound is 4-[5-(4-chlorophenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide, or a pharmaceutically-acceptable salt thereof.

5. The method of Claim 2 where the compound is 4-[5-(3-fluoro-4-methoxyphenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide, or a pharmaceutically-acceptable salt thereof.

6. The method of Claim 1 wherein the neoplasia is selected from colorectal cancer, gastrointestinal cancer, liver cancer, bladder cancer, cervical cancer, prostate cancer, lung cancer, breast cancer and skin cancer.

7. A method of preventing a neoplasia selected from adenomatous polyps, gastrointestinal cancer, liver cancer, bladder cancer, cervical cancer, prostate cancer, lung cancer, breast cancer and skin cancer, in a subject in  
 5 need of such prevention, the method comprising treating said subject with a therapeutically-effective amount of a compound of Formula II



10

wherein  $R^2$  is lower haloalkyl; wherein  $R^3$  is hydrido; and wherein  $R^4$  is phenyl optionally substituted at a substitutable position with one or more radicals selected from halo, lower alkylthio, lower alkylsulfonyl, cyano,  
 15 nitro, lower haloalkyl, lower alkyl, hydroxyl, lower alkenyl, lower hydroxyalkyl, carboxyl, lower cycloalkyl, lower alkylamino, lower dialkylamino, lower alkoxycarbonyl, aminocarbonyl, lower alkoxy, lower haloalkoxy, sulfamyl, five or six membered heterocyclic  
 20 and amino; or a pharmaceutically-acceptable salt or derivative thereof.

8. The method of Claim 7 wherein the compound is selected from compounds, and their pharmaceutically  
 25 acceptable salts, of the group consisting of  
 4-[5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
 4-[5-phenyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
 30 4-[5-(4-fluorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;

4-[5-(4-methoxyphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[5-(4-chlorophenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
5 4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[3-(difluoromethyl)-5-(4-methylphenyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[3-(difluoromethyl)-5-phenyl-1H-pyrazol-1-yl]benzenesulfonamide;  
10 4-[3-(difluoromethyl)-5-(4-methoxyphenyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[3-(difluoromethyl)-5-(3-fluoro-4-methoxyphenyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
15 4-[5-(3-fluoro-4-methoxyphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide; and  
4-[5-(4-(N,N-dimethylamino)phenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide.

20 9. The method of Claim 8 wherein the compound is 4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide, or a pharmaceutically-acceptable salt thereof.


25 10. The method of Claim 8 wherein the compound is 4-[5-(4-chlorophenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide, or a pharmaceutically-acceptable salt thereof.

30 11. The method of Claim 8 where the compound is 4-[5-(3-fluoro-4-methoxyphenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide, or a pharmaceutically-acceptable salt thereof.

35 12. A method of treating a subject suffering from a neoplastic disease state with a conjunctive therapy, said method comprising treating the subject with a

therapeutically-effective amount of a cyclooxygenase-2 selective compound and a compound selected from antibiotic-type agents, alkylating agents, antimetabolite agents, hormonal agents, immunological agents, interferon-  
 5 type agents, miscellaneous agents, metallomatrix proteases (MMP) inhibitors, SOD and  $\alpha_v\beta_3$  inhibitors.

13. The method of Claim 12 wherein the selective COX-2 inhibitor is a compound of Formula I  
 10

**I**

wherein A is a substituent selected from partially unsaturated or unsaturated heterocyclyl and  
 15 partially unsaturated or unsaturated carbocyclic rings;

wherein  $\text{R}^1$  is at least one substituent selected from heterocyclyl, cycloalkyl, cycloalkenyl and aryl, wherein  $\text{R}^1$  is optionally substituted at a  
 20 substitutable position with one or more radicals selected from alkyl, haloalkyl, cyano, carboxyl, alkoxycarbonyl, hydroxyl, hydroxyalkyl, haloalkoxy, amino, alkylamino, arylamino, nitro, alkoxyalkyl, alkylsulfinyl, halo, alkoxy and alkylthio;

25 wherein  $\text{R}^2$  is methyl or amino; and

wherein  $\text{R}^3$  is a radical selected from hydrido, halo, alkyl, alkenyl, alkynyl, oxo, cyano, carboxyl, cyanoalkyl, heterocycloxy, alkyloxy, alkylthio, alkylcarbonyl, cycloalkyl, aryl, haloalkyl,  
 30 heterocyclyl, cycloalkenyl, aralkyl, heterocyclylalkyl, acyl, alkylthioalkyl, hydroxyalkyl, alkoxycarbonyl, arylcarbonyl, aralkylcarbonyl, aralkenyl, alkoxyalkyl, arylthioalkyl, aryloxyalkyl, aralkylthioalkyl,

aralkoxyalkyl, alkoxyaralkoxyalkyl,  
alkoxycarbonylalkyl, aminocarbonyl,  
aminocarbonylalkyl, alkylaminocarbonyl, N-  
arylaminoalkyl, N-alkyl-N-arylaminoalkyl,  
5 alkylaminocarbonylalkyl, carboxyalkyl, alkylamino, N-  
arylamino, N-aralkylamino, N-alkyl-N-aralkylamino, N-  
alkyl-N-arylamino, aminoalkyl, alkylaminoalkyl, N-  
arylaminoalkyl, N-aralkylaminoalkyl, N-alkyl-N-  
aralkylaminoalkyl, N-alkyl-N-arylaminoalkyl, aryloxy,  
10 aralkoxy, arylthio, aralkylthio, alkylsulfinyl,  
alkylsulfonyl, aminosulfonyl, alkylaminosulfonyl, N-  
arylaminosulfonyl, arylsulfonyl, N-alkyl-N-  
arylaminosulfonyl; or a pharmaceutically-acceptable  
salt thereof.

15

14. The method of Claim 13 wherein A is selected  
from 5- or 6-member partially unsaturated  
heterocyclyl, 5- or 6-member unsaturated  
heterocyclyl, 9- or 10-member unsaturated condensed  
20 heterocyclyl, lower cycloalkenyl and phenyl; wherein  
R<sup>1</sup> is selected from 5- and 6-membered heterocyclyl,  
lower cycloalkyl, lower cycloalkenyl and aryl  
selected from phenyl, biphenyl and naphthyl, wherein  
R<sup>1</sup> is optionally substituted at a substitutable  
25 position with one or more radicals selected from  
lower alkyl, lower haloalkyl, cyano, carboxyl, lower  
alkoxycarbonyl, hydroxyl, lower hydroxyalkyl, lower  
haloalkoxy, amino, lower alkylamino, phenylamino,  
lower alkoxyalkyl, lower alkylsulfinyl, halo, lower  
30 alkoxy and lower alkylthio; wherein R<sup>2</sup> is methyl or  
amino; and wherein R<sup>3</sup> is a radical selected from  
hydrido, oxo, cyano, carboxyl, lower alkoxyalkyl,  
lower carboxyalkyl, lower cyanoalkyl, halo, lower  
alkyl, lower alkyloxy, lower cycloalkyl, phenyl,  
35 lower haloalkyl, 5- or 6-membered heterocyclyl, lower  
hydroxylalkyl, lower aralkyl, acyl, phenylcarbonyl,  
lower alkoxyalkyl, 5- or 6-membered heteroaryloxy,



aminocarbonyl, lower alkylaminocarbonyl, lower alkylamino, lower aminoalkyl, lower alkylaminoalkyl, phenyloxy, and lower aralkoxy; or a pharmaceutically-acceptable salt thereof.

5

15. The method of Claim 14 wherein A is selected from oxazolyl, isoxazolyl, furyl, thienyl, dihydrofuryl, pyrrolyl, pyrazolyl, thiazolyl, imidazolyl, isothiazolyl, benzofuryl, cyclopentenyl, 10 cyclopentadienyl, phenyl, and pyridyl; wherein R<sup>1</sup> is selected from pyridyl optionally substituted at a substitutable position with one or more methyl radicals, and phenyl optionally substituted at a substitutable position with one or more radicals 15 selected from methyl, ethyl, isopropyl, butyl, *tert*-butyl, isobutyl, pentyl, hexyl, fluoromethyl, difluoromethyl, trifluoromethyl, cyano, carboxyl, methoxycarbonyl, ethoxycarbonyl, hydroxyl, hydroxymethyl, trifluoromethoxy, amino, N- 20 methylamino, N,N-dimethylamino, N-ethylamino, N,N-dipropylamino, N-butylamino, N-methyl-N-ethylamino, phenylamino, methoxymethyl, methylsulfinyl, fluoro, chloro, bromo, methoxy, ethoxy, propoxy, *n*-butoxy, pentoxy, and methylthio; wherein R<sup>2</sup> is methyl or 25 amino; and wherein R<sup>3</sup> is a radical selected from hydrido, oxo, cyano, carboxyl, methoxycarbonyl, ethoxycarbonyl, carboxypropyl, carboxymethyl, carboxyethyl, cyanomethyl, fluoro, chloro, bromo, methyl, ethyl, isopropyl, butyl, *tert*-butyl, 30 isobutyl, pentyl, hexyl, difluoromethyl, trifluoromethyl, pentafluoroethyl, heptafluoropropyl, difluoroethyl, difluoropropyl, methoxy, ethoxy, propoxy, *n*-butoxy, pentoxy, cyclohexyl, phenyl, pyridyl, thienyl, thiazolyl, oxazolyl, furyl, 35 pyrazinyl, hydroxylmethyl, hydroxylpropyl, benzyl, formyl, phenylcarbonyl, methoxymethyl, furylmethyloxy, aminocarbonyl, N-

methylaminocarbonyl, N,N-dimethylaminocarbonyl, N,N-dimethylamino, N-ethylamino, N,N-dipropylamino, N-butylamino, N-methyl-N-ethylamino, aminomethyl, N,N-dimethylaminomethyl, N-methyl-N-ethylaminomethyl,  
5 benzyloxy, and phenyloxy; or a pharmaceutically-acceptable salt thereof.

16. The method of Claim 15 wherein the compound is selected from compounds, and their pharmaceutically  
10 acceptable salts, of the group consisting of

5-(4-fluorophenyl)-1-[4-(methylsulfonyl)phenyl]-3-(trifluoromethyl)pyrazole;  
4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]-1-phenyl-3-(trifluoromethyl)pyrazole;  
15 4-(5-(4-chlorophenyl)-3-(4-methoxyphenyl)-1H-pyrazol-1-yl)benzenesulfonamide  
4-(3,5-bis(4-methylphenyl)-1H-pyrazol-1-yl)benzenesulfonamide;  
20 4-(5-(4-chlorophenyl)-3-phenyl-1H-pyrazol-1-yl)benzenesulfonamide;  
4-(3,5-bis(4-methoxyphenyl)-1H-pyrazol-1-yl)benzenesulfonamide;  
4-(5-(4-chlorophenyl)-3-(4-methylphenyl)-1H-pyrazol-1-yl)benzenesulfonamide;  
25 4-(5-(4-chlorophenyl)-3-(4-nitrophenyl)-1H-pyrazol-1-yl)benzenesulfonamide;  
4-(5-(4-chlorophenyl)-3-(5-chloro-2-thienyl)-1H-pyrazol-1-yl)benzenesulfonamide;  
30 4-(4-chloro-3,5-diphenyl-1H-pyrazol-1-yl)benzenesulfonamide  
4-[5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[5-phenyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
35 4-[5-(4-fluorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;

- 4-[5-(4-methoxyphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[5-(4-chlorophenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
5 4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[4-chloro-5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[3-(difluoromethyl)-5-(4-methylphenyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
10 4-[3-(difluoromethyl)-5-phenyl-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[3-(difluoromethyl)-5-(4-methoxyphenyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
15 4-[3-cyano-5-(4-fluorophenyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[3-(difluoromethyl)-5-(3-fluoro-4-methoxyphenyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[5-(3-fluoro-4-methoxyphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
20 4-[4-chloro-5-phenyl-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[5-(4-chlorophenyl)-3-(hydroxymethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
25 4-[5-(4-(N,N-dimethylamino)phenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
5-(4-fluorophenyl)-6-[4-(methylsulfonyl)phenyl]spiro[2.4]hept-5-ene;  
4-[6-(4-fluorophenyl)spiro[2.4]hept-5-en-5-yl]benzenesulfonamide;  
30 6-(4-fluorophenyl)-7-[4-(methylsulfonyl)phenyl]spiro[3.4]oct-6-ene;  
5-(3-chloro-4-methoxyphenyl)-6-[4-(methylsulfonyl)phenyl]spiro[2.4]hept-5-ene;  
35 4-[6-(3-chloro-4-methoxyphenyl)spiro[2.4]hept-5-en-5-yl]benzenesulfonamide;

- 5-(3,5-dichloro-4-methoxyphenyl)-6-[4-(methylsulfonyl)phenyl]spiro[2.4]hept-5-ene;  
5-(3-chloro-4-fluorophenyl)-6-[4-(methylsulfonyl)phenyl]spiro[2.4]hept-5-ene;  
5 4-[6-(3,4-dichlorophenyl)spiro[2.4]hept-5-en-5-yl]benzenesulfonamide;  
2-(3-chloro-4-fluorophenyl)-4-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)thiazole;  
2-(2-chlorophenyl)-4-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)thiazole;  
10 5-(4-fluorophenyl)-4-(4-methylsulfonylphenyl)-2-methylthiazole;  
4-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)-2-trifluoromethylthiazole;  
15 4-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)-2-(2-thienyl)thiazole;  
4-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)-2-benzylaminothiazole;  
4-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)-2-(1-propylamino)thiazole;  
20 2-[(3,5-dichlorophenoxy)methyl]-4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]thiazole;  
5-(4-fluorophenyl)-4-(4-methylsulfonylphenyl)-2-trifluoromethylthiazole;  
25 1-methylsulfonyl-4-[1,1-dimethyl-4-(4-fluorophenyl)cyclopenta-2,4-dien-3-yl]benzene;  
4-[4-(4-fluorophenyl)-1,1-dimethylcyclopenta-2,4-dien-3-yl]benzenesulfonamide;  
5-(4-fluorophenyl)-6-[4-(methylsulfonyl)phenyl]spiro[2.4]hepta-4,6-diene;  
30 4-[6-(4-fluorophenyl)spiro[2.4]hepta-4,6-dien-5-yl]benzenesulfonamide;  
6-(4-fluorophenyl)-2-methoxy-5-[4-(methylsulfonyl)phenyl]-pyridine-3-carbonitrile;  
35 2-bromo-6-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]-pyridine-3-carbonitrile;  
6-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]-2-

- phenyl-pyridine-3-carbonitrile;  
4-[2-(4-methylpyridin-2-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;  
4-[2-(5-methylpyridin-3-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;  
5 4-[2-(2-methylpyridin-3-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;  
3-[1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazol-2-yl]pyridine;  
10 2-[1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazol-2-yl]pyridine;  
2-methyl-4-[1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazol-2-yl]pyridine;  
2-methyl-6-[1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazol-2-yl]pyridine;  
15 4-[2-(6-methylpyridin-3-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;  
2-(3,4-difluorophenyl)-1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazole;  
20 4-[2-(4-methylphenyl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;  
2-(4-chlorophenyl)-1-[4-(methylsulfonyl)phenyl]-4-methyl-1H-imidazole;  
2-(4-chlorophenyl)-1-[4-(methylsulfonyl)phenyl]-4-phenyl-1H-imidazole;  
25 2-(4-chlorophenyl)-4-(4-fluorophenyl)-1-[4-(methylsulfonyl)phenyl]-1H-imidazole;  
2-(3-fluoro-4-methoxyphenyl)-1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazole;  
30 1-[4-(methylsulfonyl)phenyl]-2-phenyl-4-trifluoromethyl-1H-imidazole;  
2-(4-methylphenyl)-1-[4-(methylsulfonyl)phenyl]-4-trifluoromethyl-1H-imidazole;  
35 4-[2-(3-chloro-4-methylphenyl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;  
2-(3-fluoro-5-methylphenyl)-1-[4-

- (methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazole;
- 4-[2-(3-fluoro-5-methylphenyl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;
- 5 2-(3-methylphenyl)-1-[4-(methylsulfonyl)phenyl]-4-trifluoromethyl-1H-imidazole;
- 4-[2-(3-methylphenyl)-4-trifluoromethyl-1H-imidazol-1-yl]benzenesulfonamide;
- 1-[4-(methylsulfonyl)phenyl]-2-(3-chlorophenyl)-4-trifluoromethyl-1H-imidazole;
- 10 4-[2-(3-chlorophenyl)-4-trifluoromethyl-1H-imidazol-1-yl]benzenesulfonamide;
- 4-[2-phenyl-4-trifluoromethyl-1H-imidazol-1-yl]benzenesulfonamide;
- 15 4-[2-(4-methoxy-3-chlorophenyl)-4-trifluoromethyl-1H-imidazol-1-yl]benzenesulfonamide;
- 1-allyl-4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-5-(trifluoromethyl)-1H-pyrazole;
- 20 4-[1-ethyl-4-(4-fluorophenyl)-5-(trifluoromethyl)-1H-pyrazol-3-yl]benzenesulfonamide;
- N-phenyl-[4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-5-(trifluoromethyl)-1H-pyrazol-1-yl]acetamide;
- 25 ethyl [4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-5-(trifluoromethyl)-1H-pyrazol-1-yl]acetate;
- 4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-1-(2-phenylethyl)-1H-pyrazole;
- 4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-1-(2-phenylethyl)-5-(trifluoromethyl)pyrazole;
- 30 1-ethyl-4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-5-(trifluoromethyl)-1H-pyrazole;
- 5-(4-fluorophenyl)-4-(4-methylsulfonylphenyl)-2-trifluoromethyl-1H-imidazole;
- 35 4-[4-(methylsulfonyl)phenyl]-5-(2-thiophenyl)-2-(trifluoromethyl)-1H-imidazole;

- 5- (4-fluorophenyl) -2-methoxy-4-[4-  
(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyridine;  
2-ethoxy-5- (4-fluorophenyl) -4-[4-  
(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyridine;  
5 5- (4-fluorophenyl) -4-[4- (methylsulfonyl)phenyl]-2-(2-  
propynyloxy)-6- (trifluoromethyl)pyridine;  
2-bromo-5- (4-fluorophenyl) -4-[4-  
(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyridine;  
4-[2-(3-chloro-4-methoxyphenyl)-4,5-  
10 difluorophenyl]benzenesulfonamide;  
1-(4-fluorophenyl)-2-[4-(methylsulfonyl)phenyl]benzene;  
5-difluoromethyl-4-(4-methylsulfonylphenyl)-3-  
phenylisoxazole;  
4-[3-ethyl-5-phenylisoxazol-4-yl]benzenesulfonamide;  
15 4-[5-difluoromethyl-3-phenylisoxazol-4-  
yl]benzenesulfonamide;  
4-[5-hydroxymethyl-3-phenylisoxazol-4-  
yl]benzenesulfonamide;  
4-[5-methyl-3-phenyl-isoxazol-4-yl]benzenesulfonamide;  
20 1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-  
(methylsulfonyl)benzene;  
1-[2-(4-fluoro-2-methylphenyl)cyclopenten-1-yl]-4-  
(methylsulfonyl)benzene;  
1-[2-(4-chlorophenyl)cyclopenten-1-yl]-4-  
25 (methylsulfonyl)benzene;  
1-[2-(2,4-dichlorophenyl)cyclopenten-1-yl]-4-  
(methylsulfonyl)benzene;  
1-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]-4-  
(methylsulfonyl)benzene;  
30 1-[2-(4-methylthiophenyl)cyclopenten-1-yl]-4-  
(methylsulfonyl)benzene;  
1-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-  
(methylsulfonyl)benzene;  
4-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-  
35 yl]benzenesulfonamide;  
1-[2-(4-chlorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-  
(methylsulfonyl)benzene;

- 4-[2-(4-chlorophenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;  
4-[2-(4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;  
5 4-[2-(4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;  
1-[2-(4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;  
1-[2-(2,3-difluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;  
10 4-[2-(3-fluoro-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;  
1-[2-(3-chloro-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;  
15 4-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;  
4-[2-(2-methylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;  
ethyl 2-[4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]oxazol-2-yl]-2-benzyl-acetate;  
20 2-[4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]oxazol-2-yl]acetic acid;  
2-(tert-butyl)-4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]oxazole;  
25 4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]-2-phenyloxazole;  
4-(4-fluorophenyl)-2-methyl-5-[4-(methylsulfonyl)phenyl]oxazole; and  
4-[5-(3-fluoro-4-methoxyphenyl)-2-trifluoromethyl-4-oxazolyl]benzenesulfonamide.  
30

17. The method of Claim 16 wherein the compound is selected from compounds, and their pharmaceutically acceptable salts, of the group consisting of  
35 4-[5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;



4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
4-[5-(3-fluoro-4-methoxyphenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;  
5 3-[1-[4-(methylsulfonyl)phenyl]-4-trifluoromethyl-1H-imidazol-2-yl]pyridine;  
2-methyl-5-[1-[4-(methylsulfonyl)phenyl]-4-trifluoromethyl-1H-imidazol-2-yl]pyridine;  
4-[2-(5-methylpyridin-3-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;  
10 4-[5-methyl-3-phenylisoxazol-4-yl]benzenesulfonamide;  
4-[5-hydroxymethyl-3-phenylisoxazol-4-yl]benzenesulfonamide;  
[2-trifluoromethyl-5-(3,4-difluorophenyl)-4-oxazolyl]benzenesulfonamide;  
15 4-[2-methyl-4-phenyl-5-oxazolyl]benzenesulfonamide;  
and  
4-[5-(3-fluoro-4-methoxyphenyl-2-trifluoromethyl)-4-oxazolyl]benzenesulfonamide.

20

18. The method of Claim 16 wherein the compound is 4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide, or a pharmaceutically-acceptable salt thereof.

25

19. The method of Claim 1 wherein the neoplasia is adenomatous polyps.

20. The method of Claim 7 wherein the neoplasia is adenomatous polyps.

30

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 97/18670

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K31/415 A61K31/10 A61K31/18

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	WO 96 38418 A (G.D. SEARLE & CO.) 5 December 1996 see claims 1-3,9-16 see page 4, line 35 - page 5, line 1 ---	1-20
Y	WO 95 15316 A (G. D. SEARLE & CO.) 8 June 1995 cited in the application see claims 16-18,34-36 see page 8, line 4 - line 16 ---	1-20
Y	B. S. TEICHER ET AL: "Cyclooxygenase and lipoxigenase inhibitors as modulators of cancer therapies" CANCER CHEMOTHERAPY AND PHARMACOLOGY, vol. 33, 1994, pages 515-522, XP000676574 see the whole document ---	1-20
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

29 January 1998

Date of mailing of the international search report

13. 03. 98

Name and mailing address of the ISA  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Siatou, E

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/18670

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>HUANG P ET AL: "CYCLOOXYGENASE AND 5-LIPOXYGENASE INHIBITORS FOR THE PREVENTION AND TREATMENT OF CANCER" EXPERT OPINION ON INVESTIGATIONAL DRUGS, vol. 4, no. 3, March 1995, pages 243-249, XP000603398 see the whole document ---</p>	1-20
A	<p>WO 95 15318 A (G.D. SEARLE &amp; CO.) 8 June 1995 cited in the application see the whole document ---</p>	1-20
A	<p>G. ARA ET AL: "Cyclooxygenase and lipoxigenase inhibitors in cancer therapy" PROSTAGLANDINS, LEUKOTRIENES AND ESSENTIAL FATTY ACIDS, vol. 54, 1996, pages 3-16, XP002053643 ---</p>	1-20
A	<p>D. L. EARNEST ET AL: "Piroxicam and Other Cyclooxygenase Inhibitors: Potential for Cancer Chemoprevention" JOURNAL OF CELLULAR BIOCHEMISTRY, vol. Suppl. 16I, 1992, pages 156-166, XP002053644 see abstract ---</p>	1-20
A	<p>C. MILLIGAN-CIHA ET AL: "Inhibition of tumor growth by intratumor administration of cyclooxygenase inhibitors" FEDERATION PROCEEDINGS, vol. 42, no. 3, 1 March 1983, USA, page 682 XP002053645 Abstract No. 2285 see abstract -----</p>	1-20

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 97/18670

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
**see FURTHER INFORMATION sheet PCT/ISA/210**
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

**FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

Claims Nos.: 1-20

because they relate to subject matter not required to be searched by this Authority, namely:

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

Remark : Although claims 1-20 are directed to a method of treatment of the human/animal body , the search has been carried out and based on the alleged effects of the compound/composition.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No

PCT/US 97/18670

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9638418 A	05-12-96	AU 5886296 A	18-12-96
-----			
WO 9515316 A	08-06-95	US 5466823 A	14-11-95
		US 5521207 A	28-05-96
		AU 1171495 A	19-06-95
		CA 2177576 A	08-06-95
		CN 1141630 A	29-01-97
		CZ 9601503 A	11-12-96
		EP 0731795 A	18-09-96
		FI 962249 A	29-05-96
		HU 74180 A	28-11-96
		JP 9506350 T	24-06-97
		NO 962184 A	29-05-96
		PL 314695 A	16-09-96
		US 5510496 A	23-04-96
		US 5563165 A	08-10-96
		US 5508426 A	16-04-96
		US 5516907 A	14-05-96
		US 5504215 A	02-04-96
		ZA 9409418 A	28-11-95
-----			
WO 9515318 A	08-06-95	US 5434178 A	18-07-95
		AU 1171595 A	19-06-95
		CA 2177574 A	08-06-95
		EP 0731796 A	18-09-96
		JP 9505830 T	10-06-97
-----			